NEW!
KILLS OVER 250 WEED TYPES
AS LISTED, INCLUDING CRABGRASS

GUARANTEED RESULTS†

Spectracide®

WEED STOP®

FOR LAWNS

plus Crabgrass Killer

HERBICIDA PARA CÉSPEDES Y CONTRA MANO DE CANGREJO

- VISIBLE RESULTS IN 8 HOURS
- WON'T HARM LAWNS*
- KILLS OVER 250 WEED TYPES

AS LISTED, INCLUDING DANDELION, CLOVER, CRABGRASS & YELLOW NUTSEDGE

†For terms of guarantee, see back panel.

ACTIVE INGREDIENTS:	
2,4-D, dimethylamine salt	0.253%
Quinclorac	0.121%
Dicamba, dimethylamine salt	0.029%
Sulfentrazone	0.015%
INERT INGREDIENTS:	99.582%
TOTAL	100.000%

THIS PRODUCT CONTAINS:
0.018 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 0.210%
0.010 lb. 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 0.121%
0.002 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 0.024%
0.001 lb. N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide per gallon or 0.015%
Isomer Specific by AOAC Method.

KEEP OUT OF REACH OF CHILDREN
CAUTION See back booklet for additional Precautionary Statements.

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCIÓN Consulte el folleto en la parte posterior para conocer las precauciones adicionales.

Net 1 gal (128 fl oz / 3.7 L)

READY TO USE™

17-12371

Spectracide

WEED STOP®

FOR LAWNS

plus Crabgrass Killer

HERBICIDA PARA CÉSPEDES Y CONTRA MANO DE CANGREJO

PRODUCT FACTS

WHAT IT DOES:

Kills major broadleaf and troublesome grass weeds as listed, including dandelion, chickweed and clover as well as crabgrass, foxtail, and yellow nutsedge. (See inside for complete list)

WHEN TO USE:

Apply when daytime temperatures are between 45°F and 90°F

WHERE TO USE:

Lawns: Kills the weeds not the lawn

QUESTIONS & COMMENTS:

Call 1-800-917-5438 or visit our website at www.spectracide.com

†GUARANTEED RESULTS

GUARANTEED RESULTS OR YOUR MONEY BACK

If you are not satisfied with this product for any reason, send your written request for a refund to the address below, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

NOTA: El comprador asume toda la responsabilidad en caso de no seguir las instrucciones sobre seguridad y modo de empleo

MADE IN THE USA

Distributed by Spectrum Group
Div. of United Industries Corp.
PO Box 142642, St. Louis, MO 63114-0642

PROOF OF PURCHASE

FPO UPC
0 71121 95663 7

EPA Reg. No. 2217-887-8845 EPA Est. Nos. 39754-WI-1 (R), 9688-MO-1 (U), 79533-WI-1 (BT), 39609-MO-1 (BM) Circled letter is first letter of lot number. 17-12370 © 2007 UIC

ABRA AQUÍ **PEEL HERE**

STOP **READ THE ENTIRE LABEL FIRST.**
ALTO. LEA TODA LA ETIQUETA ANTES DE USAR.

WEEDS CONTROLLED/MALEZAS CONTROLADAS
BROADLEAF WEEDS CONTROLLED/MALEZAS DE HOJA ANCHA CONTROLADAS:
Amaranth, purple; Amaranth, slender; Aster, heath; Aster, purple; Aster, white heath; Aster, white prairie; Artichoke; Austrian fieldcress; Bedstraw, catchweed; Bedstraw, smooth; Bedstraw, northern; Bedstraw, pineywoods; Beggarticks, hairy; Betony, Florida; Bindweed, field; Bindweed, hedge; Bitter wintercress; Bittercress, hairy; Bittersweed; Black-eyed Susan; Black medic; Blood flower milkweed; Blue lettuce; Brassbuttons; Bristly oxtongue; Broomweed; Buckhorn; Bullnettle; Burclover; Burdock; Burweed, lawn (spurweed); Buttercup, bulbous; Buttercup, creeping; Buttonweed, common (poorjoe); Buttonweed, Virginia; Carolina geranium (wild); Carpetweed; Catnip; Catsear, spotted (false dandelion); Chamber bitter; Chickweed, common; Chickweed, mousear; Chickweed, sticky; Chicory; Cinquefoil; Clover, al sike; Clover, crimson; Clover, hop; Clover, rabbit foot; Clover, red; Clover, strawberry; Clover, yellow sweet; Clover, white sweet; Clover, white; Cockle; Cocklebur; Cocklebur, oriental; Cocklebur, spiny; Common mullein; Corn spurry; Creeping beggarweed; Creeping jenny (creeping charlie); Cupid shaving brush; Cudweed; Daisy, English; Daisy, oxeye; Dandelion, common; Dandelion, false; Day flower; Deadnettle; Dichondra; Dock, broadleaf; Dock, curly; Dogbane; Dogfennel; Dollarweed (pennywort); Doveweed; Elderberry; Falseflax; False sunflower; Fiddleneck; Filaree, redstem; Filaree, whitestem; Fleabane, daisy (annual); Fleabane, rough; Galinsoga, hairy; Galinsoga, smallflower; Goldenrod; Ground ivy; Groundsel; Gumweed; Hairy fleabane; Hawkweed, meadow; Hawkweed, mousear; Hawkweed, orange; Hawkweed, yellow; Healall; Heartleaf drymary; Hemp; Henbit; Hoary cress; Horsenettle; Horseweed; Jimsonweed; Johnny-jumpup violet; Knawel; Knotweed, prostrate; Kochia; Lambsquarters; Lespedeza; Mallow, alkali; Mallow, bristly; Mallow, common; Mallow, dwarf; Mallow, Venice; Marcela; Matchweed; Mexicanweed; Moneywort; Morningglory, bigroot; Morningglory, ivyleaf; Morningglory, Mexican; Morningglory, red; Morningglory, tall; Morningglory, woolly; Mugwort; Mustard, ball; Mustard, black; Mustard, blue; Mustard, hedge; Mustard, Indian; Mustard, tansy; Mustard, tumble; Mustard, white; Mustard, wormseed; Nettle, stinging (burning); Nettle, tall; Nettle, wood; Parsley-piert; Parsnip; Pearlwort; Pennycress (Frenchweed); Pepperweed, perennial; Pepperweed, Virginia (peppergrass); Pigweed, prostate; Pigweed, redroot; Pigweed, Russian; Pigweed, smooth; Pigweed, tumble; Pineappleweed; Plains coreopsis (tickseed); Plantain, blackseed; Plantain, bracted; Plantain, broadleaf; Plantain, buckhorn; Plantain, hoary; Plantain, narrowleaf; Plantain, slender; Plantain, woolly; Poison ivy; Poison oak; Pokeweed; Prairie sunflower; Prickly lettuce (compass plant); Prickly sida; Prostrate knotweed; Puncturevine; Purslane, common; Pusley, Brazil; Pusley, Florida; Pusley, large flower; Ragweed, bur; Ragweed, common; Ragweed, lanceleaf; Ragweed, western; Redstem filaree; Rough cinquefoil; Scarlet pimpernel; Shepherdspurse; Smartweed, ladysthumb; Smartweed, pale; Smartweed, Pennsylvania; Smooth chaff-flower; Smooth dock; Sorrel, red (sheep); Sowthistle, annual; Sowthistle, spiny; Spanish needles; Speedwell, bi-lobed; Speedwell, common; Speedwell, corn; Speedwell, ivyleaf; Speedwell, Persian; Speedwell, purslane; Speedwell, slender; Speedwell, snow; Speedwell, thymeleaf; Spiny amaranth; Spurge, spotted; Spurge, prostrate; Spurweed; Star of Bethlehem; Strawberry, Indian mock; Tansy ragwort; Tanweed; Thistle, blessed; Thistle, bull; Thistle, Canada; Thistle, flodman (prairie); Thistle, musk; Thistle, platte; Thistle, plumeless; Thistle, Russian; Thistle, Scotch; Thistle, wavyleaf; Thistle, yellow (horrible); Trailing crownvetch; Velvetleaf; Vervain, blue; Vervain, hoary; Vervain, prostrate; Vervain, tall; Vetch, bird; Vetch, hairy; Vetch, milk; Vetch, narrowleaf; Vetch, wild; Virginia creeper; Western clematis; Western salsify; Wild aster; Wild buckwheat; Wild carrot; Wild four-o-clock; Wild garlic; Wild lettuce; Wild marigold; Wild mustard (yellow rocket); Wild onion; Wild parsnip; Wild radish; Wild rape; Wild strawberry; Wild sweet potato; Wild violet; Woodsorrel, creeping; Woodsorrel, violet; Woodsorrel, yellow (oxalis); Woolly croton; Wormseed; Yarrow, common; Yarrow, western; Yarrow, yellow

GRASSY WEEDS/MALEZAS HERBOSAS: Crabgrass, large and smooth; Foxtail, green, giant and yellow; Yellow nutsedge

STOP **READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

General Precautions and Restrictions:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

For home lawn care:

Spectracide® Weed Stop® for Lawns plus Crabgrass Killer is a ready-to-use (RTU) product intended for "spot" treatment of individual broadleaf and grass weeds by homeowners on residential lawns. This product should be applied to lawns containing the following species of grass: Kentucky bluegrass, perennial ryegrass, fescue species (including tall, red, and fine leaf varieties), Bermudagrass*, buffalograss and zoysiagrass.

*Application to Bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to Bermudagrass when daytime temperatures exceed 85°F.

This product should **not** be used on lawns containing bentgrass, St. Augustinegrass, bahiagrass, centipedegrass, or carpetgrass. Do **not** use on dichondra or lawns with desirable clovers or legumes, on vegetable gardens, fruits, or ornamentals (flowers, trees, groundcovers, hedges, landscape/flower beds and shrubs).

- This 1 gal retail container size treats approximately 1,000 weeds or up to 1,000 sq ft.

HAND SPRAYER PRODUCT

1. Shake well.
2. Connect trigger sprayer. Unscrew and discard the white cap on the bottle. Insert the hose into the bottle and tighten the cap securely.
3. Prime sprayer: Turn the sprayer nozzle one-half turn counterclockwise. Hold the sprayer lower than the bottle and squeeze the trigger several times until it sprays. Adjust the nozzle to coarse spray pattern to minimize wind drift.
4. Spray just enough Spectracide Weed Stop for Lawns plus Crabgrass Killer to lightly cover broadleaf and grass weeds. ***Excessive spraying can injure some lawns.*** Do not apply when daytime temperatures are above 90°F as lawn injury can occur. Spray only the weeds, not the entire lawn.
5. Weeds will begin wilting and browning overnight.

USE TIPS

- Some hard-to-kill weeds may require re-treatment. Wait at least 3 weeks between applications — some weeds can take that long to die.
- Rainproof — rain or watering 3 hours after application will not wash away effectiveness.
- Do not allow spray to drift onto desirable plants since injury may result.

ALTO **LEA PRIMERO TODA LA ETIQUETA. CUMPLA TODAS LAS PRECAUCIONES Y SIGA CUIDADOSAMENTE LAS INSTRUCCIONES.**

MODO DE EMPLEO

Se considera una violación a la ley federal usar este producto con otros fines que los indicados en la etiqueta.

Precauciones y restricciones generales:

- No aplique este producto de modo que entre en contacto con personas o mascotas, ya sea directamente o por arrastre. Mantenga a las personas y las mascotas fuera del área durante la aplicación.
- No permita que las personas ni las mascotas entren en el área tratada hasta que los rocíos se hayan secado.

Para el cuidado del césped residencial:

El herbicida para céspedes y contra mano de cangrejo Spectracide® Weed Stop® es un producto listo para usar, fabricado para el tratamiento "localizado" de pastos y malezas de hoja ancha, por parte de propietarios, en céspedes residenciales. Este producto debería aplicarse en céspedes que contengan las siguientes especies de pasto: césped azul de Kentucky, ryegrass perenne, especies de festuca (incluyendo las variedades de hoja alta, roja y fina), césped Bermuda* césped búfalo y césped zoysia.

*La aplicación al césped Bermuda puede provocar que el césped se vuelva amarillo o se decolore temporalmente pero se puede esperar una recuperación completa. No lo aplique al césped Bermuda cuando las temperaturas diurnas superan los 85°F.

Este producto **no** se debe usar en céspedes que contengan agróstide, césped San Agustín, césped bahía, césped ciempiés o césped alfombra. **No** lo use en céspedes dichondra o con tréboles o legumbres deseables, en huertas de verduras o frutas, ni en jardines con plantas ornamentales (flores, árboles, cubresuelos, setos, lechos de flores o plantas y arbustos).

- Este tamaño de envase de venta de 1 galón trata aproximadamente 1,000 malezas o hasta 1,000 pies cuadrados.

PRODUCTO CON ROCIADOR DE GATILLO

1. Agite bien.
2. Conecte el gatillo del rociador. Desenrosque y deseche la tapa de la botella. Introduzca la manguera en la botella y ajuste la tapa de modo que quede segura.
3. Cebe el rociador: Gire la boquilla del rociador media vuelta en sentido contrario al de las manecillas del reloj. Sostenga el rociador debajo del nivel de la botella y apriete el gatillo varias veces hasta que salga el rocío. Ajuste la boquilla para descargar un rocío grueso para minimizar el arrastre por el viento.
4. Rocíe el herbicida para céspedes y contra mano de cangrejo Spectracide Weed Stop en cantidad suficiente para cubrir ligeramente el césped y las malezas de hoja ancha. ***El exceso de rocío puede dañar algunos tipos de césped.*** No aplique cuando las temperaturas diurnas sean superiores a 90°F ya que se puede dañar el césped. Rocíe sólo las malezas, no todo el césped.
5. Las malezas comenzarán a marchitarse y adquirir color marrón durante la noche.

CONSEJOS SOBRE EL USO

- Algunas malezas difíciles de eliminar pueden requerir la repetición del tratamiento. Deje pasar un período de al menos 3 semanas entre las aplicaciones; algunas malezas pueden demorar todo ese tiempo en morir.
- A prueba de lluvia. La lluvia o el riego después de 3 horas de la aplicación no reducen la eficacia.
- No permita el arrastre del rocío sobre plantas deseables ya que puede causarles daño.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Turn the nozzle to "OFF" position and store with sprayer above the level of liquid in container to prevent leakage. Store in original container in an area inaccessible to children and pets.

DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards To Humans & Domestic Animals:

CAUTION: Causes moderate eye injury. Avoid contact with eyes or clothing. Wear protective eyewear. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves. It is recommended that safety glasses include front, brow and temple protection. Wash thoroughly with soap and water after handling. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

First Aid	
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes:	• Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-917-5438 for emergency medical treatment advice.	

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Questions & Comments? Call 1-800-917-5438 or visit our website at www.spectracide.com

ALMACENAMIENTO Y ELIMINACIÓN

ALMACENAMIENTO DEL PESTICIDA: Gire la boquilla hasta la posición "OFF" (apagada) y guárdelo con el rociador por encima del nivel de líquido en el envase para evitar derrames. Guarde en su envase original, en un lugar que no sea accesible ni a los niños ni a las mascotas.

ELIMINACIÓN:

Si está vacío: No vuelva a usar este envase. Tírelo en la basura, u ofrézcalo para reciclar, si existe esa opción.

Si está parcialmente vacío: Llame a su servicio de eliminación de desechos más cercano para obtener instrucciones de eliminación. Nunca tire el producto restante por ningún drenaje interno ni externo.

PRECAUCIONES

Riesgos para los seres humanos y los animales domésticos:

PRECAUCIÓN: Causa lesión moderada a los ojos. Evite el contacto con los ojos o la ropa. Use gafas de protección. Nocivo si se ingiere. Cuando utilice este producto, use camisa de manga larga, pantalones largos, calcetines, zapatos y guantes de goma/hule. Se recomienda que las gafas de protección incluyan protección delantera para la frente y las sienes.

Después de manipular el producto, lávese bien con agua y jabón. Después de utilizar este producto, enjuague los guantes antes de quitárselos, quítese la ropa y lávela por separado antes de volver a usarla y rápidamente lávese bien las manos y la piel que estuvo expuesta, con agua y jabón. Quítese la ropa mojada y báñese en regadera tan pronto como sea posible.

Primeros Auxilios	
Si se ingiere:	• Llame inmediatamente a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento. • Si la persona puede tragar, haga que beba poco a poco un vaso de agua. • No induzca el vómito a menos que así lo indique el centro de control de envenenamientos o el médico. • No administre nada por la boca a una persona que haya perdido el conocimiento.
Si entra en contacto con los ojos:	• Mantenga los ojos abiertos y enjuague con agua lentamente, con cuidado, durante 15-20 minutos. Si hay lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. • Llame a un centro de control de envenenamientos para recibir instrucciones de tratamiento.
Cuando llame a un centro de control de envenenamientos o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto. También puede llamar al 1-800-917-5438 para obtener información acerca del tratamiento médico de urgencia.	

RIESGOS PARA EL MEDIO AMBIENTE:

Este pesticida es tóxico para los peces y los invertebrados acuáticos. No aplique directamente al agua. El arrastre y el escurrimiento pueden resultar peligrosos para los organismos acuáticos de las zonas cercanas al sitio de aplicación.

NOTA: El comprador asume toda la responsabilidad en caso de no seguir las instrucciones sobre seguridad y modo de empleo.

¿Preguntas o comentarios? Llame al 1-800-917-5438.

MADE IN THE USA Distributed by Spectrum Group, Division of United Industries Corporation, PO Box 142642, St. Louis, MO 63114-0642 EPA Reg. No. 2217-887-8845
EPA Est. Nos. 39754-WI-1 Ⓡ, 9688-MO-1 Ⓤ, 79533-WI-1 (BT), 39609-MO-1 (BM) Circled letter is first letter of lot number. 17-12370 ©2007 UIC